

Attractor Neural Networks

Giorgio Parisi
Dipartimento di Fisica
Università di Roma *La Sapienza*

February 1, 2008

Abstract

In this lecture I will present some models of neural networks that have been developed in the recent years. The aim is to construct neural networks which work as associative memories. Different attractors of the network will be identified as different internal representations of different objects. At the end of the lecture I will present a comparison among the theoretical results and some of the experiments done on real mammal brains.

1 Introduction

The field of neural networks has been intensively studied by many people coming from quite different disciplines: mathematics, computer sciences, physics, biology and psychology. Each of them carries with him his training, the methodology of his discipline and the taste for what he believes it is important or interesting.

The goals of all these people are quite different. Roughly speaking they can be divided into those who want to understand how the real mammal brain works and those who want to use the information they can gather in the study of the brain in order to build new brands of computers. These new computers should behave in a more intelligent way, with more good sense compared to the existing ones.

There are also others who are interested in understanding how complex collective behaviour emerges from the interaction of many simpler elementary objects. This is quite a general issue. Everywhere in biology it can be observed that an assembly of objects behaves quite differently from the isolated objects and this is a crucial phenomenon. Examples of emergent collective behaviour are macromolecules from atoms, cells from macromolecules, pluricellular animals from single cells, species, evolution and ecosystems from animals.

The task of deducing the laws for the higher levels from those of the lower level is not simple at all. Many physicists (and myself in particular [1]) are fascinated by this last question, also because in the past we have already successfully answered to similar questions, albeit in a much simpler context. The emergence of collective behaviour is quite familiar to us under the name of phase transition. Many examples can be found: an assembly of spins becomes ferromagnetic decreasing the temperature (or increasing it, if we follow the original notation by Celsius), a system of many electrons behaves as superconductor...

Ferromagnetism, superconductivity are collective phenomena that can be understood only using the appropriate tools of statistical mechanics. For many systems, especially for superconductivity, the understanding of the behaviour of the whole system from the microscopic components and the construction of the corresponding theoretical framework has not been an easy job. Much more elaborate constructions are needed in order to understand some peculiar behaviour of disordered systems, like spin glasses, where many microscopically different, but macroscopically equivalent equilibrium states are present [2], [3].

Nowadays physicists have succeeded in reaching a good control of many of the observed collective phenomena in physical systems and I believe that the task of deriving macroscopic laws from the microscopic ones in biological setting is our duty. It seems to me that it is necessary to use the techniques of statistical mechanics in order to reach an understanding of the dynamics and of the mutual interaction

(also with DNA) of the three thousands different enzymes in an *Escherichia Coli*. The times seem to be ripe for this task also because in the last ten years we have started to understand the behaviour of physical complex systems, mainly disordered systems as spin glasses, which display many different equilibrium states. Our toolbox of statistical mechanics contains rather sharp and sophisticated instruments.

It approach to biology physicists have often reproached for oversimplifying. I believe that physicists do oversimplify (using the standards of biologists) and they do it for good reasons. Let me present the two most important reasons:

- Modern physics start at Galileo's times with the seemly unnatural choice of neglecting friction. In every day life friction dominates; the force is proportional to the speed as in the old approach of Aristotle, not to the acceleration. Without friction we are unable even to walk, as it can be verified when we find ice on the road. A world without friction would not look like the real world.

However physics started by first extrapolating from the experiments the properties of a frictionless world and by constructing a theoretical framework able to describe this artificial world ¹, which culminated in Newtons Mechanics. Friction has been included in the theory at a much later stage.

Many times in the history of physics progress have been achieved by selecting some features that are supposed to be simpler to analyse and by neglecting the rest for the time being, up to the moment in which the theory was able to cope with new features and nowadays this approach is an automatic reflex of physicists. The great simplifications (which we will see later) introduced by Hopfield in its study of Attractor Neural Networks are a magnificent step backward, in the tradition of physics.

- There is also a more technical reason for oversimplifying, which has been discovered in the study of second order phase transitions by Kadanoff and Wilson. It carries the name of *universality*. The basic idea is simple. The number of qualitatively different collective behaviours of a system at the macroscopic level is not very high and it is definitively much less than the number of different microscopic systems. Thousands (or millions) of different ferromagnetic materials (of Ising type) exist and in all of them the specific heat diverges at the phase transition point as $|T - T_c|^\alpha$. The value of α (about -1.1) is the same for all the systems. The qualitative and quantitative properties of

¹Let me quote from the introduction from a book of Tartaglia (freely translated into modern language): *In this book we will study those objects that move with constant speed in the horizontal direction and with constant acceleration in the vertical direction. If real cannon balls do not follow this law, the worse for them, we will not speak about them.*

the collective behaviour depend on the kind of collective behaviour that is observed, not on the microscopic details of the system.

Of course quite different system may have qualitative different phase transitions (some become ferromagnetic, others superfluids or glasses). Phase transitions may be divided in a few universality classes (at least for homogeneous systems)²; the qualitative behaviour and some of the quantitative features near the phase transition point are constant inside a given universality class and they do not change with small change of the systems. In other words the collective behaviour can be understood in self consistent way without making reference to the microscopic dynamics. This program reached its higher success for two dimensional systems, where all possible collective behaviour of systems belonging to a quite large class has been classified using simple consistency requirements for the collective behaviour.

If the understanding of the collective behaviour can be obtained by studying the universality classes and not a particular system, quite different systems that are supposed to belong to the same universality class are equivalent as far as the collective behaviour is concerned. One of the first reaction of a theoretical physicist to a new problem is to search the simplest system that should belong to the same universality class. Universality suggests that unwanted complications can be neglected and often we do not need to reintroduce them at a later stage, because they do not change the main results. Of course a large amount of experience and maybe of Serendipity is needed to discriminated the unwanted complications from the essential properties.

In this lecture I will describe some of the developments than have been done in the recent years on the construction of a network of neurons which should behave as associative memory. Associative memory is memory in the usual human sense³: we remember names, faces, notions, we recognise known objects and we are able to recall them also if we receive only a partial information . At the end of the lecture I will present a comparison of the predictions of the models with experiments with real mammal brains.

²In the same spirit of Thom's classification of catastrophes

³The word associative is used to distinguish this kind of memory by the computer memory (addressable memory) where each address identifies the content of the address.

2 A Schematic Description of Neurons

Before entering into the details of the model it may be convenient to present a schematic description of the neurons and their mutual interactions[4]. We will mainly consider the electric activity of the neuron, also because it is the easiest to measure. If we consider the difference in voltage of the neuron with the surrounding liquid, we observe fast spikes of about 100 mV of amplitude of a time duration of about a millisecond. The number of spikes per second may range from 1 to a few hundreds, although in normal physiological conditions is not greater than a few tens.

The activity of a neuron depends on the signal it receives from other neurons. Indeed different neurons are connected by synapses, which are unidirectional devices needed to transmit information. If the neuron A has a synapse toward B , when A is active, neurotransmitters are released by A and absorbed by B . The release of neurotransmitters increases when the activity of the neuron increases.

The synapses may be excitatory or inhibitory. The neurotransmitters increase the activity of B if the synapse is excitatory, and they decrease the activity if the synapse is inhibitory. A given neuron may receive signals from many neurons (the number of neurons is 10^4 in average, with a range which goes from 2 to 10^5). The activity of a neuron will be high if the excitatory signals overcame the inhibitory ones, otherwise it would be low.

Different synapses will have different effects. The efficacy of a given synapse depends on time and it is usually believed that there is some physiological mechanism which strengthens those synapses in which both neurons are simultaneously active (the Hebb's rule). The details of this mechanism in vertebrates are unclear at the present moment, although a large amount of information has been collected.

Synapses allow a neuron to send a message specifically to another neuron, which may be quite far away. Neurons also send non specific messages to all nearby neurons by producing neurotransmitters which diffuse in the intercellular liquid. Here we are not going to consider this activity.

If we try to construct a model for a set of N interconnected neurons (i.e. a neural network), we can parametrize the activity of the neurons by N functions $\sigma_i(t)$, $i = 1, \dots, N$, and the synaptic strength by N^2 functions $J_{i,k}(t)$.

The stimulus of the network on a given neuron (i) is assumed to be given by

$$S_i(t) = \sum_{k=1,N} J_{i,k}(t)\sigma_k(t), \quad (1)$$

i.e. the total stimulus is simply the sum of the stimulus coming from each neuron, which we assume to be equal to the product of the synaptic strength with the activity

of the neuron producing the stimulus ⁴.

The dynamic equations for the neuron are supposed to be in the simplest case:

$$\frac{d\sigma_i}{dt} = f(\sigma_i, S_i), \quad (2)$$

where f is a non linear function of its arguments.

In order to fully describe the network, expecially in the learning stage, we should also give the equations which control the time evolution of the synaptic strengths (J), however for reasons of time I will not discuss this point, which is very interesting and unfortunately not so well understood. In the following we will assume that the dynamics of the J 's has been such to produce synaptic couplings with of the kind we postulate.

3 Associative memory

It is believed that an internal representation (i.e. a pattern of neuron activities) is associated to each object or category that we are able to recognise and remember. It is also believed that an object is memorised by changing the synaptic strengths in an appropriate way. In order to have an associative memory the dynamics of the neurons should such that

- If the neurons are set equal to μ_i at time 0, where the μ_i correspond to one of the learned pattern, the neuron activities $\sigma_i(t)$ should remain for all times not too far from μ_i . In other words the learned pattern correspond to a stable attractor points, in a statistical sense (i.e. small fluctuations around the fixed points are allowed).
- If we apply an external stimulus which push the neurons in the direction of being equal to τ_i , and if the τ_i only partially correspond to one of the learned patterns μ_i , the neurons should evolve toward the μ_i . This property implies that the memory is able to retrieve the information on the whole object from the knowledge of a part of it or in presence of partially wrong information.
- If the τ_i of the external stimulus are very different from the μ_i of any stored pattern, the system should recognised the absence of a match. This goal may be obtained if in this case no fixed point is reached and a chaotic behaviour is observed. In this case the memory is called cognitive because it knows

⁴In writing eq. (1) we have neglected the propagation time of the stimulus. Adding a variability in the propagation times leads to the appearance of new phenomena.

when a correct result is obtained: reaching an attractor (a fixed point) has the meaning of recalling the information, while no stable point implies the absence of recognition.

It is clear that physiological constraint implies that many different patterns of activities may be stored same neural network. Our model of network should be such that many different attractors are present. From this point of view memory is very similar to a spin glass, where many different equilibrium states are possible[2, 3].

4 The Hopfield model

The Hopfield model is the simplest version of a model in which many different attractors do exist [5]. In this model the neuron activities are conventionally taken to be ± 1 and the time t is also an integer valued quantity. This choice is rather unnatural; the choice $\sigma = 0$ or 1 , would sound much better, however no crucial qualitatively difference are among these two cases, as we shall see later.

One step (i.e. the increase of t by an unity) for the dynamics of the neuron activities is obtained by applying (for all i) the rule

$$\sigma_i(t + \frac{i+1}{N}) = \text{sign}(S_i(t + \frac{i}{N})), \quad (3)$$

where we still have

$$S_i(t) = \sum_{k=1, N} J_{i,k} \sigma_k(t). \quad (4)$$

If the matrix J is symmetric ($J_{i,k} = J_{k,i}$), this dynamics corresponds to a sequential algorithm for looking for the minimum of the Hamiltonian

$$H = - \sum_{i,k=1, N} J_{i,k} \sigma_i \sigma_k. \quad (5)$$

The main advantage of the Hopfield model was that at this stage the model is very similar to the zero temperature dynamics of a statistical mechanics model. The analogy becomes more strong is we introduce the possibility of errors and we write

$$\begin{aligned} \sigma_i(t + \frac{i+1}{N}) &= \text{sign}(S_i(t + \frac{i}{N})) \text{ with probability } |\tanh(S_i((t + \frac{i}{N})/T))|, \\ \sigma_i(t + \frac{i+1}{N}) &= -\text{sign}(S_i(t + \frac{i}{N})) \text{ with probability } 1 - |\tanh(S_i((t + \frac{i}{N})/T))|. \end{aligned} \quad (6)$$

In this case we obtain the dynamics of a statistical mechanics system at temperature T with the random Hamiltonian (5), which can successfully studied using the

techniques developed for spin glasses. In the limit $T \rightarrow 0$, we recover the original error free dynamics of eq. (3).

It is possible to prove that for the original zero temperature dynamics a fixed point is reached for large time, i.e. the σ 's become time independent. Indeed the Hamiltonian decreases every times that a σ changes its value. The absence of limiting cycles is a very strong simplification and it is the main reason for considering this model ⁵.

Our aim is to store in network $M = \alpha N$ different patterns, which are denoted by μ_i^p , where the index i runs from 1 to N and the index p runs from 1 to M . The matrix J is assumed to have the following symmetric form

$$J_{i,k} = \sum_{p=1,M} \mu_i^p \mu_k^p, \quad (7)$$

which has a strong Hebbian flavour. Indeed it correspond to say that the matrix J changes by the quantity $\mu_i^p \mu_k^p$ each time that the network learns the new pattern μ_i^p . The symmetry of the matrix J , although is not physiologically realist, is necessary for using the statistical mechanics techniques and for being able to consider the evolution of the network as a process of energy minimisation.

We can ask the question if there are configurations of the σ which are fixed point of the dynamics, i.e. they satisfy the equations

$$\sigma_i = \text{sign}(\sum_{k=1,N} J_{i,k} \sigma_k) \quad (8)$$

and are not far from one of the μ^p 's for some value of p .

This requirement may be stated in a more precise way. We can define the distance among a configuration σ and a configuration τ as

$$d(\sigma, \tau) = 1/N \sum_{i=1,N} (\sigma_i - \tau_i)^2 / 4. \quad (9)$$

In other words $d(\sigma, \tau)$ is the percentage of neurons which are different in the two configurations. If the two configurations are uncorrelated, the distance is .5.

We look for existence of configurations σ , which satisfy eq. (8) and have a small distance with one of the μ 's. This problem has been carefully studied in the limit where N goes to infinity[6]. In the simplest case we assume that the different μ 's

⁵We could also allow the possibility of having a periodic behaviour with period 2, i.e. the σ becomes time independent separately for odd and even times. In this case we should update the neurons in a sequential way, i.e. by setting $\sigma_i(t+1) = \text{sign}(S_i(t))$. The properties of this model are rather similar to the one described in the text and it is possible that only fixed points or limiting cycles of period two may be present.

are uncorrelated one from the other and they have in average the same number of +1 and -1. In this situation one finds the existence of two regimes:

- For $\alpha < \alpha_c$, where $\alpha_c \approx .14$, one can find at least a configuration σ very near to each of the M configurations μ^p . More precisely we can find it at a distance less than $d_{Max}(\alpha)$, where $d_{Max}(\alpha)$ is an increasing function of α , which reaches its maximum (i.e. 0.01) in this interval at α_c . The number of these solutions is increasing with N , but very slowly (i.e. as $\exp(\omega N)$ with a very small value of ω).
- For $\alpha > \alpha_c$ no solutions exist with small distance and one find solution only at distances of order of $.4 - .5$.

In the first regime the network works as an associative memory, however, when the loading become greater than α_c , the memory stops to work and goes into a state of total confusion. This first order transition is a feature of the model which disappear changing some of the characteristics of the model. In all models there exist a critical value of α such that the system does not work as a memory for $\alpha > \alpha_c$.

5 More refined models

The Hopfield model contains many unrealistic features. Many more refined models have been proposed. The restriction to the neuron activity to be ± 1 can be easily removed without dramatic consequences. Different choices of the probability distribution of the patterns and of the learning procedure have been studied. Other features can be added, e.g. the dilution of the network, which consists in assuming that not all the neurons are connected and some of the J 's are strictly zero.

I cannot mention these interesting results for lack of time. I will only recall a very interesting result obtained by the late Elisabeth Gardner [7]. Here we do not commit ourselves to any learning algorithm and we can ask if we can find a matrix J such that all the patterns are fixed point of the dynamics. A very interesting computation show that when the patterns are uncorrelated, in the limit $N \rightarrow \infty$ this is always possible for $\alpha < 1$ and never possible for $\alpha > 1$. The interest in this kind of results is that they tell us what a network can do independently of learning algorithm, i.e. of the dynamics for the J s.

Using a symmetric form for the matrix J we remain in the framework of equilibrium statistical mechanics. New qualitative features appear if the J are asymmetric [8]. Indeed in the Hopfield model the evolution of the σ_i is such that a fixed point is always reached, also if this fixed point has nothing to do with one of the stored patterns. If we introduce asymmetries in the matrix J , it can be shown that the

evolution does not leads always to a fixed point and the spurious fixed points not correlated with the stored patterns do disappear. If the network does not reach a fixed point near one of the stored pattern, the time evolution of the network is chaotic and the variables σ_i will change value in an apparent random way. This results is very interesting because one obtains without effort a good feature (the possibility of chaotic states) by removing one of the artificial characteristic of the model (i.e. the symmetry of the J 's).

A very simple model in which this effect can be studied is the asymmetrically diluted model, where

$$J_{i,k} = J_{i,k}^H A_{i,k}, \quad (10)$$

$J_{i,k}^H$ being given by eq.(7) and $A_{i,k}$ is an asymmetric matrix, whose elements are 1 with probability z and 0 with probability $1 - z$.

In the limit of small z the model can be easily studied analytically and after simple computations one finds that the storage capacity α_c is equal to $2z/\pi$ and that the large time behaviour of the network is fully chaotic when it is not near one of the attractors [9].

A very large amount of work has been done in these recent years also on many other problems (e.g. learning of rules from example) [10, 11, 12, 13] and in most of the cases the extension of the techniques developed for statistical mechanical systems has been very fruitful.

6 Real experiments

We have seen that we can construct many model with different features which work as associative memories. It is quite likely that these results will be eventually important for the projects of new generations of computers. At the present moment a very interesting practical application is to use neural networks as fast triggers in order in high energy experiments. i.e. at super colliders. In this case one should decide if an event is interesting or not in a rather short time, of the order of 10 nanoseconds [14]. Here a neural network may be able to perform this rather complex task in such a short time using its massive parallelism, i.e. using many neurons, each of them being physically implemented in Silicium.

From a biological point of view we are interested to know if attractors do really exist in mammal brains and how the brain perform very complex task (for example it is able to recognise if different image do correspond to the same physical object). In order to address to this problem we have to identified experimentally the memory cell in which the information is stored.

In the past many neurons have been found which respond (i.e. they switch from a low firing rate to an higher one) in a selective way to a given external stimulus (for example visual). The stimulus was sometimes highly complex (e.g. a face), sometimes very simple (an horizontal line). Unfortunately up to a recent time the activity of the neurons disappeared when the stimulus was removed.

Very recently neurons have been found which remain active in a selective way after the removal of the stimulus for a period larger than 16 seconds [15, 16]. In the experiments monkeys, of the species *Macaca Refus*, were trained to learn and to memorise 100 fractal patterns, which were presented in a given order.

After learning these patterns were presented to the monkeys in pairs at a time distance of about 20 seconds. They were supposed to act differently if the first pattern was equal to the first and a fruit juice was given to them if they performed correctly. It was found that in small area of about 1 mm^3 , there are neurons which respond in selective way to the visual presentation of only few of the patterns.

It is reasonable to suppose that the recorded activities come from cells that belong to the attractor, not from cells that are excited by the attractor. If this hypothesis is correct, it should be possible to start an experimental study of the property of the attractor. Many interesting features of the experiment have been compared with theoretical predications.

A very interesting and unexpected feature of the experiment is the existence of correlations in the neural activities of different patterns. The patterns have been presented in a fixed, randomly chosen order. The patterns itself are uncorrelated, however the probability that a given neuron is active after the presentation of the i^{th} pattern is much higher than the average, if the neuron is active after the presentation of k^{th} pattern and $|i - k|$ is not large (e.g. smaller than 6). The neural activities corresponding to patterns which have been learned at nearby times are correlated.

It has been shown [17] that these correlations arise naturally in a relative simple model of neural network and that the experimental features of the correlations is in good agreement with the theoretical predictions.

It is quite likely that in the future experiments and theory will evolve in a convergent way and it will be possible to obtain a much more detailed confirmation (or refutation) of the theoretical approach presented here.

I hope to have given to the audience some flavour of the large amount of work that has been done in the field of neural network in the last years. Much more could have been said, but the time is over.

References

- [1] G. Parisi, Physics Word, (1993).
- [2] M. Mézard, G. Parisi and M.A. Virasoro, *Spin glass theory and beyond*, World Scientific, Singapore (1987).
- [3] G. Parisi, *Field Theory, Disorder and Simulations*, World Scientific, Singapore (1992).
- [4] V. Braitenberg and A. Shulz, *Anatomy of the cortex*, Springer-Verlag, Berlin (1991).
- [5] J.J. Hopfield, Proc. Natl. Acad. Sci. USA 79, 2554 (1982).
- [6] D.J. Amit, *Modeling Brain Functions*, Cambridge, Cambridge University Press (1989).
- [7] E. Gardner J. Phys. A, 22, 76 (1987).
- [8] G. Parisi J. Phys. A, 19, L675 (1986).
- [9] B. Derrida, E. Gardner and A. Zippelius, Europhys. Lett. 4, 167 (1987).
- [10] J. Denker, D. Schwartz, B. Wittner, S. Solla, R. Horward, L. Jackel and J.J. Hopfield, Complex Systems, 1, 877 (1987).
- [11] H. Somplinsky and N. Tishby, Phys Rev Letters 65, 1683 (1990).
- [12] G. Parisi, Network 3, ?? (1992).
- [13] See for example Oppen preprint ????
- [14] *Neural Networks: from biology to high energy physics* O. Benhar, C. Bosio, P. Del Giudice, E. Tabet eds., Pisa, ETS Editrice (1991) .
- [15] Y. Miyashita and Chang H.S., Nature, 331, 68 (1988).
- [16] Y. Miyashita, Nature, 335, 817 (1988).
- [17] D.J. Amit, N. Brunel and M.V. Tsodyks *Correlations of cortical Hebbian reverberations: experiment and theory*, Rome I preprint N.964 (1993).